

Analysis of inflationary cosmological models in gauge theories of gravitation

A V Minkevich^{1,2} and A S Garkun¹

¹ Department of Theoretical Physics, Belarussian State University, F. Skoriny av. 4, Minsk 20050, Belarus

² Department of Physics and Computer Methods, Warmia and Mazury University in Olsztyn, Poland

E-mail: minkav@bsu.by, garkun@bsu.by

Abstract. Inflationary homogeneous isotropic cosmological models filled by scalar fields and ultrarelativistic matter are examined in the framework of gauge theories of gravitation. By using quadratic scalar field potential numerical analysis of flat, open and closed models is carried out. Properties of cosmological models are investigated in dependence on indefinite parameter of cosmological equations and initial conditions at a bounce. Fulfilled analysis demonstrates regular character of all cosmological models.

PACS numbers: 04.50.+h; 98.80.Cq; 11.15.-q

1. Introduction

As it was shown in a number of papers (see [1, 2] and Refs. given therein), general feature of homogeneous isotropic cosmological models built in the framework of gauge theories of gravitation (GTG) [3, 4] is their regular bouncing character. All cosmological solutions of generalized cosmological Friedmann equations (GCFE) deduced for homogeneous isotropic models in GTG are regular in metrics, Hubble parameter and its time derivative because of gravitational repulsion effect at extreme conditions (extremely high energy densities and pressures), which takes place in the case of usual gravitating systems satisfying energy dominance condition [5]. In the case of inflationary cosmological models filled by interacting scalar field and usual gravitating matter, indicated above property of cosmological solutions was obtained without concretization explicit form of scalar field potential and equation of state of gravitating matter [1]. However, we have to use their explicit form to obtain and analyze cosmological solutions of GCFE.

The present paper is devoted to analysis of inflationary cosmological models filled by noninteracting scalar fields and ultrarelativistic matter. In Sec. 2 the GCFE for such inflationary cosmological models are introduced. In Sec. 3 some general properties of its solutions are examined. In Sec. 4 and 5 by using quadratic scalar field potential numerical analysis of cosmological solutions for flat, open and closed models is carried out.

2. Generalized cosmological Friedmann equations in GTG

The GCFE deduced in the frame of Poincare GTG and metric-affine GTG have the following form [6-8]

$$\frac{k}{R^2} + \left\{ \frac{d}{dt} \ln \left[R \sqrt{|1 + \alpha(\rho - 3p)|} \right] \right\}^2 = \frac{8\pi G}{3} \frac{\rho + \frac{\alpha}{4}(\rho - 3p)^2}{1 + \alpha(\rho - 3p)}, \quad (1)$$

$$\frac{\left[\dot{R} + R \left(\ln \sqrt{|1 + \alpha(\rho - 3p)|} \right)' \right]}{R} = -\frac{4\pi G}{3} \frac{\rho + 3p - \frac{\alpha}{2}(\rho - 3p)^2}{1 + \alpha(\rho - 3p)}, \quad (2)$$

where $R(t)$ is the scale factor of Robertson-Walker metrics, ρ is energy density, p is pressure, $k = +1, 0, -1$ for closed, flat, open models respectively, G is Newton's gravitational constant (the light velocity $c = 1$), α is indefinite parameter with inverse dimension of energy density, a dot denotes differentiation with respect to time t . Together with Newton's gravitational constant G the parameter α is fundamental physical constant of relativistic isotropic cosmology built in the frame of GTG. The value of α^{-1} determines the scale of extremely high energy densities [5], which will be supposed to be less than the Planckian one. From equations (1)–(2) follows the energy conservation law in usual form

$$\dot{\rho} + 3H(\rho + p) = 0 \quad \left(H = \frac{\dot{R}}{R} \right). \quad (3)$$

Obviously, the solutions of GCFE (1)–(2) depend on values of G and α . However, we can exclude them and transform equations (1)–(2) to dimensionless form by means of the following transformation of the time t and functions $R(t)$, $\rho(t)$, $p(t)$

$$\begin{aligned} t &\rightarrow \tilde{t} = t \sqrt{\frac{G}{\alpha}}, & R &\rightarrow \tilde{R} = R \sqrt{\frac{G}{\alpha}}, \\ \rho &\rightarrow \tilde{\rho} = \alpha \rho, & p &\rightarrow \tilde{p} = \alpha p \end{aligned} \quad (4)$$

As result we obtain

$$\frac{k}{\tilde{R}^2} + \left[\frac{d}{d\tilde{t}} \ln \left(\tilde{R} \sqrt{|1 + \tilde{\rho} - 3\tilde{p}|} \right) \right]^2 = \frac{8\pi}{3} \frac{\tilde{\rho} + \frac{1}{4}(\tilde{\rho} - 3\tilde{p})^2}{1 + \tilde{\rho} - 3\tilde{p}}, \quad (5)$$

$$\frac{\left[\tilde{R}' + \tilde{R} \left(\ln \sqrt{|1 + \tilde{\rho} - 3\tilde{p}|} \right)' \right]}{\tilde{R}} = -\frac{4\pi}{3} \frac{\tilde{\rho} + 3\tilde{p} - \frac{1}{2}(\tilde{\rho} - 3\tilde{p})^2}{1 + \tilde{\rho} - 3\tilde{p}}, \quad (6)$$

where a prime denotes the differentiation with respect to $\tilde{t}$. The form of equation (3) does not change by transformations (4)

$$\tilde{\rho}' + 3\tilde{H}(\tilde{\rho} + \tilde{p}) = 0 \quad \left(\tilde{H} = \frac{\tilde{R}'}{\tilde{R}}, \quad \tilde{H} = H \sqrt{\frac{\alpha}{G}} \right). \quad (7)$$

In the case of models filled by scalar field ϕ with potential $V = V(\phi)$ and gravitating matter with energy density ρ_m and pressure p_m we have

$$\rho = \frac{1}{2}\dot{\phi}^2 + V + \rho_m, \quad p = \frac{1}{2}\dot{\phi}^2 - V + p_m. \quad (8)$$

By using the transformation (4) we obtain from (8) the following dimensionless expressions:

$$\tilde{\rho} = \frac{1}{2}\tilde{\phi}'^2 + \tilde{V} + \tilde{\rho}_m, \quad \tilde{p} = \frac{1}{2}\tilde{\phi}'^2 - \tilde{V} + \tilde{p}_m, \quad (9)$$

where $\tilde{\phi} = \phi\sqrt{G}$, $\tilde{V} = \alpha V$, $\tilde{\rho}_m = \alpha \rho_m$, $\tilde{p}_m = \alpha p_m$. By neglecting the interaction between scalar field and gravitating matter, from conservation law (7) follows the equation for scalar field

$$\tilde{\phi}'' + 3\tilde{H}\tilde{\phi}' = -\frac{\partial\tilde{V}}{\partial\tilde{\phi}} \quad (10)$$

and conservation law for gravitating matter

$$\tilde{\rho}_m' + 3\tilde{H}(\tilde{\rho}_m + \tilde{p}_m) = 0. \quad (11)$$

To integrate equations (5)–(6) with $\tilde{\rho}$ and $\tilde{p}$ defined by (9) we have to know the equation of state of gravitating matter $p_m = p_m(\rho_m)$. To build realistic cosmological models we have to take into account the change of equation of state under their evolution in accordance with properties of gravitating matter, but it is generally a complicated physical problem. To study cosmological models near a bounce and simplify further analysis we will use below the equation of state for ultrarelativistic matter $p_m = \frac{1}{3}\rho_m$.

In considered case from equation (11) follows the integral for gravitating matter

$$\tilde{\rho}\tilde{R}^4 = \text{const}. \quad (12)$$

Then the GCFE (5)–(6) can be transformed in accordance with Ref. [5] to the following form:

$$\left[\tilde{H} \left(\tilde{Z} + 3\tilde{\phi}'^2 \right) + 3\frac{\partial\tilde{V}}{\partial\tilde{\phi}}\tilde{\phi}' \right]^2 + \frac{k}{\tilde{R}^2} \tilde{Z}^2 = \frac{8\pi}{3} \left[\tilde{\rho}_m + \frac{1}{2}\tilde{\phi}'^2 + \tilde{V} + \frac{1}{4} \left(4\tilde{V} - \tilde{\phi}'^2 \right)^2 \right] \tilde{Z}, \quad (13)$$

$$\begin{aligned} \tilde{H}' \left(\tilde{Z} + 3\tilde{\phi}'^2 \right) + 3\tilde{H}^2 \left(\tilde{Z} - \tilde{\phi}'^2 \right) + 3 \left[\frac{\partial^2\tilde{V}}{\partial\tilde{\phi}^2}\tilde{\phi}'^2 - \left(\frac{\partial\tilde{V}}{\partial\tilde{\phi}} \right)^2 \right] \\ = 8\pi \left[\tilde{V} + \frac{1}{3}\tilde{\rho}_m + \frac{1}{4} \left(4\tilde{V} - \tilde{\phi}'^2 \right)^2 \right] - \frac{2k}{\tilde{R}^2} \tilde{Z}, \end{aligned} \quad (14)$$

where $\tilde{Z} = 1 + 4\tilde{V} - \tilde{\phi}'^2$.

The greatest part of solutions of GCFE (13)–(14) are inflationary cosmological solutions having direct physical interest. Properties of these solutions are studied below in sections 3–5 in dependence on initial conditions at a bounce, restrictions on parameter α and parameters of scalar field potential. Note that, if the value of α^{-1} is much less than the Planckian energy density, equations (13)–(14) with scalar field potentials applying in inflationary cosmology lead also to oscillating solutions studied in Ref. [9].

3. Some properties of inflationary cosmological solutions

Let us introduce 4-dimensional phase space P with the axes $(\tilde{\phi}, \tilde{\phi}', \tilde{H}, \tilde{\rho}_m)$. Then some trajectory in space P corresponds to any solution of GCFE. Trajectories of cosmological solutions are situated in area of P , where $\tilde{Z} \geq 0$ and, hence, are limited by the bounds L defined as

$$\tilde{Z} = 0 \quad \text{or} \quad \tilde{\phi}' = \pm \sqrt{1 + 4\tilde{V}}. \quad (15)$$

According to equation (13) values of the Hubble parameter are equal to

$$\tilde{H}_{\pm} = \frac{-3 \frac{\partial \tilde{V}}{\partial \tilde{\phi}} \tilde{\phi}' \pm \sqrt{\tilde{D}}}{\tilde{Z} + 3\tilde{\phi}'^2}, \quad (16)$$

where

$$\tilde{D} = \frac{8\pi}{3} \left[\tilde{\rho}_m + \frac{1}{2}\tilde{\phi}'^2 + \tilde{V} + \frac{1}{4} \left(4\tilde{V} - \tilde{\phi}'^2 \right)^2 \right] \tilde{Z} - \frac{k}{\tilde{R}^2} \tilde{Z}^2. \quad (17)$$

Two values of the Hubble parameter $\tilde{H}_+$ and $\tilde{H}_-$ correspond to two different solutions named as $\tilde{H}_{\pm}$ -solutions. In points of bounds L we have $\tilde{D} = 0$, $\tilde{H}_+ = \tilde{H}_-$ and the Hubble parameter is determined by

$$\tilde{H}_L = -\frac{1}{\tilde{\phi}'} \frac{\partial \tilde{V}}{\partial \tilde{\phi}}. \quad (18)$$

Now let us consider extremal H_0 -surfaces in the space P , in points of which $\tilde{H}_+ = 0$ or $\tilde{H}_- = 0$. Denoting the values of quantities on H_0 -surfaces by means of index “0”, we obtain from (13) the following equation for H_0 -surfaces, which are in fact in 3-dimensional hyperspace $\tilde{H} = 0$ of the phase space P :

$$\frac{k}{\tilde{R}_0^2} \tilde{Z}_0^2 + 9 \left(\frac{\partial \tilde{V}}{\partial \tilde{\phi}} \right)_0^2 \tilde{\phi}_0'^2 = \frac{8\pi}{3} \left[\tilde{\rho}_{m0} + \frac{1}{2}\tilde{\phi}_0'^2 + \tilde{V}_0 + \frac{1}{4} \left(4\tilde{V}_0 - \tilde{\phi}_0'^2 \right)^2 \right] \tilde{Z}_0, \quad (19)$$

where $\tilde{Z}_0 = 1 + 4\tilde{V}_0 - \tilde{\phi}_0'^2$. For given scalar field potential $\tilde{V}(\tilde{\phi})$ this equation determines H_0 -surfaces for flat models ($k = 0$), and in the case of closed and open models equation (19) gives parametric families of H_0 -surfaces with parameter $\tilde{R}_0$. According to equation (14) the time derivative of the Hubble parameter $\tilde{H}'_0$ on H_0 -surfaces is

$$\begin{aligned} \tilde{H}'_0 = & \left\{ 8\pi \left[\tilde{V}_0 + \frac{1}{3}\tilde{\rho}_{m0} + \frac{1}{4} \left(4\tilde{V}_0 - \tilde{\phi}_0'^2 \right)^2 \right] \right. \\ & \left. - 3 \left[\left(\frac{\partial^2 \tilde{V}}{\partial \tilde{\phi}^2} \right)_0 \tilde{\phi}_0'^2 - \left(\frac{\partial \tilde{V}}{\partial \tilde{\phi}} \right)_0^2 \right] - \frac{2k}{\tilde{R}_0^2} \tilde{Z}_0 \right\} \left(\tilde{Z}_0 + 3\tilde{\phi}_0'^2 \right)^{-1}. \end{aligned} \quad (20)$$

If $\tilde{H}'_0 > 0$, we have in corresponding points of H_0 -surfaces the transition from compression to expansion (a bounce), and in the case $\tilde{H}'_0 < 0$ the transition from expansion to compression takes place. Equations (19)–(20) permit to find specific points of solutions, where $\tilde{H} = 0$ and $\tilde{H}' = 0$ simultaneously.

To analyze solutions of GCFE we have to study H_0 -surfaces defined by (19) for given scalar field potential. We will use below the simplest quadratic potential in the form $V = \frac{1}{2}m^2\phi^2$ or $\tilde{V} = \frac{1}{2}\tilde{m}^2\tilde{\phi}^2$, where $\tilde{m} = m\sqrt{\frac{\alpha}{G}}$. In this case H_0 -surfaces depend on parameter $\tilde{m}$ and the value of scale factor $\tilde{R}_0$. In the case of $\tilde{\rho}_m = 0$ equation (19) for H_0 -surfaces includes only variables $\tilde{\phi}_0$ and $\tilde{\phi}'_0$, and the analysis of H_0 -surfaces as well as bounds L is reduced to analysis of corresponding curves on the plane $(\tilde{\phi}, \tilde{\phi}')$. Numerical analysis will be given below in the Planckian system of units with $\hbar = c = G = 1$. Note, that admissible values of dimensionless potential $\tilde{V} = \alpha V$ can be essentially greater than 1, if the scale of extremely high energy densities defined by α^{-1} is much less then the Planckian one. Hence, values of parameter $\tilde{m}$ can be large. Below the analysis is given for various types of cosmological inflationary models.

4. Numerical analysis of flat inflationary models

In the case of flat models ($k = 0$) H_0 -curves $B_{1,2}$ are presented in Fig. 1 together with bounds $L_{1,2}$ having the form of two hyperbolas $\tilde{\phi}' = \pm\sqrt{1 + 2\tilde{m}^2\tilde{\phi}^2}$. Each of two curves $B_{1,2}$ contains two parts corresponding to vanishing of $\tilde{H}_+$ or $\tilde{H}_-$ and denoting by (B_{1+}, B_{2+}) and (B_{1-}, B_{2-}) respectively [1]. The parts (B_{1+}, B_{1-}) and (B_{2+}, B_{2-}) have the common points $(\tilde{\phi} = 0, \tilde{\phi}' = \pm 1)$ with the bounds L_1 and L_2 on the axis $\tilde{\phi}'$. From equations (19) – (20) it is possible to show, that if

$$\tilde{m} < 2\sqrt{\frac{\pi}{42}}\sqrt{-3 - 47\left(1009 + 84\sqrt{159}\right)^{-1/3} + \left(1009 + 84\sqrt{159}\right)^{1/3}} \approx 1.346,$$

than the value of $\tilde{H}'_0$ on H_0 -curves is positive (this means that $\tilde{H}'_{+0} > 0$ in points of B_{1+} and B_{2+} , and $\tilde{H}'_{-0} > 0$ in points of B_{1-} and B_{2-}). In this case $B_{1,2}$ -curves are “bounce curves” [1], in points of which the transition from compression to expansion takes place. If $\tilde{m} > 1.346$ (that corresponds to large values of α), the regions with $\tilde{H}'_0 < 0$ appear on H_0 -curves near the axis $\tilde{\phi}'$ for sufficiently small values of $|\tilde{\phi}|$ (see Fig. 1). In this case, besides regular cosmological solutions, the GCFE have scalar field oscillating solutions [9].

Now let us analyze properties of cosmological solutions for flat models. Each such solution, generally speaking, includes the quasi-de-Sitter compression and inflationary stages, stage after inflation (Fig. 2) and the stage of transition from compression to expansion (Fig. 3). Properties of cosmological solutions — duration of different stages of evolution, characteristics of scalar field oscillations and behaviour of the Hubble parameter after inflation and before the compression stage are studied in dependence on initial conditions at a bounce and the value of $\tilde{m}$. Like to GR [10], the initial value of $\tilde{\phi}_0$ must be greater than 1 ($\tilde{\phi}_0 > 1$) to appear inflationary stage.

Because GCFE lead to slow-rolling regime which coincide practically with that of GR, properties of inflationary stage of our models are the same as in GR. Namely, the duration of inflationary stage t_{infl} increases linearly by increasing of $|\tilde{\phi}_0|$ and decreases inversely proportional by increasing of $\tilde{m}$ (Fig. 4).




Figure 1. Bound $L_{1,2}$ -curves and bounce curves $B_{1,2}$ for flat models in the case of potential $\tilde{V} = \frac{1}{2}\tilde{m}^2\tilde{\phi}^2$. Dotted curves correspond to negative values of $\tilde{H}'_0$.




Figure 2. Compression and inflationary stages; postinflationary stage.




Figure 3. Transition stage




Figure 4. Duration of inflationary stage in dependence on values $\tilde{\phi}_0$ and $\tilde{m}$.

Because of $\tilde{\phi}'_0 \neq 0$, cosmological solutions are asymmetric in the time with respect to the bounce point. In particular, the duration of quasi-de-Sitter stage of compression is not equal to duration of inflationary stage. In the case of bounce in points of curves B_{1+} and B_{2+} the duration of inflationary stage is greater than duration of quasi-de-Sitter compression stage. If the bounce takes place in points of bounce curves B_{1-} and B_{2-} the situation is opposite.

For given value of $\tilde{m}$ inflationary models without quasi-de-Sitter compression stage ($\tilde{\rho}_m = 0$) can be realized for certain interval of $\tilde{\phi}_0$. For example, at $\tilde{m} = 1.2$ corresponding interval of $\tilde{\phi}_0$ is (28, 100), then the growth of the scale factor $R(t)$ is in interval (e^{130}, e^{1450}) . By increasing of $\tilde{m}$ the upper and lower bounds of interval for $\tilde{\phi}_0$ of discussed solutions increase linearly (Fig. 5a). In the case of presence of radiation ($\tilde{\rho}_m \neq 0$) the lower bound of interval of $\tilde{\phi}_0$ for solutions without quasi-de-Sitter stage of compression decreases essentially and upper bound decreases slowly (Fig. 5b).

The most principal feature of our cosmological models is the presence of regular transition stage from compression to expansion. The behaviour of scalar field $\tilde{\phi}$ and the Hubble parameter $\tilde{H}$ at transition stage is essentially nonlinear and complicated. The duration of transition $\Delta\tilde{t}_{\text{tr}}$ stage decreases by increasing of each of parameters $\tilde{m}$, $\tilde{\phi}_0$ and $\tilde{\rho}_{m0}$ (Fig. 6 – Fig. 7).

After inflation stage the scalar field oscillates near its minimum. These oscillations don't depend on initial values of $\tilde{\phi}_0$ and $\tilde{\rho}_{m0}$, but essentially depend on parameter $\tilde{m}$. Like to GR frequency and amplitude of scalar field oscillations decreases with the time, but quantitatively such dependence is different. Note, that energy density $\tilde{\rho}_m$ is negligibly small at the end of inflationary stage, even if $\tilde{\rho}_{m0} \neq 0$. Application of Krylov-Bogoliubov method [11] of solutions of the system of differential equations (10) and (14) with $\tilde{\rho}_m = 0$, after some tedious calculations, leads to the following asymptotics for scalar field $\tilde{\phi}(t)$ and the scale factor $\tilde{R}(t)$ at late time

$$\tilde{\phi}(\tilde{t}) \approx \frac{1}{\sqrt{3\pi \tilde{m}\tilde{t}}} \cos\left(\frac{3\tilde{m}^2 + 2\pi}{8\pi\tilde{m}\tilde{t}} + \tilde{m}\tilde{t}\right), \quad (21)$$




Figure 5. The upper and lower bounds of interval for admissible values of $\tilde{\phi}_0$ for solutions without quasi-de-Sitter compression stage.




Figure 6. Duration of transition stage in dependence on values of $\tilde{m}$ and $\tilde{\phi}_0$.

$$\tilde{R}(\tilde{t}) \approx \sqrt[3]{16\pi} \tilde{R}_0 \tilde{t}^{2/3} - \frac{\sqrt[3]{16\pi} \tilde{R}_0}{\tilde{m}^2 \tilde{t}^{4/3}} \cos \left(\frac{27\tilde{m}^4 + 36\pi\tilde{m}^2 - 20\pi^2}{12\pi\tilde{m}(3\tilde{m}^2 + 2\pi)} \tilde{t} + 2\tilde{m}\tilde{t} \right). \quad (22)$$

Equation (21) shows, that the amplitude of scalar field oscillations $\tilde{A}$ decreases with time as follows $\tilde{A} = \tilde{a}(\tilde{m})\tilde{t}^{-1}$, where $\tilde{a} = 4\tilde{m}^{-1}/\sqrt{3}$. Numerical analysis confirms relations (21)–(22) at sufficiently small values of $\tilde{m}$ ($\tilde{m} \lesssim 30$), in particular, the amplitude of oscillations decreases inverse proportional to time (at $\tilde{m} = \text{const}$) and inverse proportional to $\tilde{m}$ (Fig. 8).[‡] But at large values of $\tilde{m}$ ($\tilde{m} \sim 1000$) conclusions obtained by means of Krylov-Bogoliubov method differ from numerical results. In particular, the inverse proportional dependence between the amplitude of scalar field oscillations and the value of $\tilde{m}$ is not valid (Fig. 9). This is connected with the fact, that in the case of

[‡] According to numerical calculation, constant of proportionality at Fig. 8b is equal to 3.070, which is close to $\sqrt{3\pi} \approx 3.069$ as Krylov-Bogoliubov method requires.




Figure 7. Duration of transition stage in dependence on $\tilde{\rho}_{m0}$.

large values of $\tilde{m}$ (i.e. large values of α) the Hubble parameter, unlike to GR, oscillates near the value $\tilde{H} = 0$ (Fig. 10). The frequency of $\tilde{H}$ -oscillations is two times greater than the frequency of $\tilde{\phi}$ -oscillations. The value of $\tilde{H}$ becomes positive with the time and tends asymptotically to zero like to GR. Note that, there are similar $\tilde{\phi}$ -oscillations and $\tilde{H}$ -oscillations before the compression stage also.




Figure 8. Dependence of amplitude of $\tilde{\phi}$ -oscillations on the time $\tilde{t}$ and on $\tilde{m}$ at $\tilde{m} < 30$.

5. Numerical analysis of open and closed models

In the case of open models H_0 -curves are similar to that of flat models and the family of H_0 -curves is situated on the plan $(\tilde{\phi}, \tilde{\phi}')$ between curves L_1 and B_1 , and also between curves L_2 and B_2 . At $R_0 \rightarrow 0$ H_0 -curves of open models tend to bounds $L_{1,2}$, and at $R_0 \rightarrow \infty$ they approach to curves $B_{1,2}$ (see Fig. 1). In accordance with expression (21) for $\tilde{H}'_0$ the minimum value of $\tilde{m}$, by which negative values of $\tilde{H}'_0$ appear for open models, depends on $\tilde{R}_0$ and it is smaller then that for flat models. Properties of cosmological solutions for open models are similar to that of flat models. There are area of $\tilde{H}' < 0$




Figure 9. Dependence of $\tilde{\phi}$ -oscillations amplitude at large values of $\tilde{m}$.




Figure 10. $\tilde{H}$ -oscillations after inflation at large values of $\tilde{m}$.

on H_0 -curves, if $\tilde{m} > 2\sqrt{\pi/6} \approx 1.45$ and the value of $\tilde{R}_0$ is arbitrary or $\tilde{m} < 2\sqrt{\pi/6}$ and

$$\tilde{R} < \sqrt{\frac{3}{4\pi}} \sqrt{\frac{1 + 24\tilde{m}^2}{1 + 30\tilde{m}^2 + 216\tilde{m}^4}}.$$

Note that in the case of open models there are also solutions with $\tilde{Z} < 0$, which are separated from cosmological solutions by bounds L_1 and L_2 . These solutions are hardly ever have physical sense.

The form of H_0 -curves for closed models ($k = +1$) differs essentially from that for flat and open models, and it depends on the value of $\tilde{R}_0$ for given value of $\tilde{m}$. For sufficiently large values of $\tilde{R}_0$ H_0 -curves for closed models include besides two curves similar to that of flat model also one closed curve with center in origin of coordinates (Fig. 5). If $\tilde{m} < 1.34$ and the value of $\tilde{R}_0$ is sufficiently large, the value of $\tilde{H}'_0$ is positive on H_0 -curves (a) and on the part of H_0 -curve (b). In the case $\tilde{m} > 1.34$ the region with $\tilde{H}'_0 < 0$ appears on H_0 -curves (a) near the axes $\tilde{\phi}'$. By decreasing of $\tilde{R}_0$ the H_0 -curves of closed models are deformed and transformed into H_0 -curves presented in Fig. 12. If $\tilde{m} < 1.4$ H_0 -curves (c) contain regions with $\tilde{H}'_0 > 0$ and regions with $\tilde{H}'_0 < 0$; small region with $\tilde{H}'_0 < 0$ can be present on H_0 -curves (d) although on the greatest part of H_0 -curves (d) we have $\tilde{H}'_0 > 0$. In the case $\tilde{m} > 1.4$ and $\tilde{R}_0 < \tilde{R}_{01}$ (for example, $\tilde{R}_{01} = 1.5$ for $\tilde{m} = 2.1$, $\tilde{R}_{01} = 2.5$ for $\tilde{m} = 3.5$)§ we have $\tilde{H}'_0 > 0$ on the curves (f) and $\tilde{H}'_0 < 0$ on the H_0 -curves (e). Like to flat and open models oscillating solutions exist for large values $\tilde{m}$ (α). Unlike to flat and open models, in the case of closed models there are cosmological solutions symmetric in the time with respect to a bounce ($\tilde{\phi}'_0 = 0$). Note, that cosmological solutions have properties similar to flat models, if the value of $\tilde{R}_0$ at a bounce is sufficiently large.

§ The value $\tilde{R}_{01}^2$ is the real root of cubic equation, which can be obtained from equation (19) $\pi(-243\tilde{m}^6 - 306\pi\tilde{m}^4 - 408\pi^2\tilde{m}^2 + 32\pi^3)\tilde{R}_{01}^6 + 3(6561\tilde{m}^8 + 7776\pi\tilde{m}^6 + 12960\pi^2\tilde{m}^4 - 5376\pi^3\tilde{m}^2 + 256\pi^4)\tilde{R}_{01}^4 - 6912\tilde{m}^2\pi(81\tilde{m}^4 - 144\pi\tilde{m}^2 + 16\pi^2)\tilde{R}_{01}^2 + 3981312\tilde{m}^4\pi^2 = 0$




Figure 11. $\tilde{H}_0$ -curves for closed models at large values $\tilde{R}_0$.




Figure 12. $\tilde{H}_0$ -curves for closed models at small values of $\tilde{R}_0$.

6. Conclusion

Numerical analysis of inflationary cosmological models carried out in the framework of gauge theories of gravitation confirms obtained earlier conclusion about their regular character. The difference in behaviour of cosmological models at the end of inflationary stage in comparison with GR (in the case of large values of α) can be interesting,

if we take into account recent results concerning the anisotropy of cosmic microwave background [12]. This means that further investigations of perturbations in discussed inflationary models is of direct physical interest.

- [1] Minkevich A V 2005 (*Preprint* gr-qc/0506140), submitted to *Gravitation & Cosmology*
- [2] Minkevich A V 2005 *Int. J. Mod. Phys. A* **20** 2436–42.
- [3] Hehl F W 1980 *In: Cosmology and Gravitation* (New York: Plenum Press)
- [4] Hehl F W, McGrea G D, Mielke E W and Neeman Y 1995 *Phys. Rep.* **258** 1
- [5] Minkevich A V 2005 Gravitational repulsion effect at extreme conditions (*Preprint* gr-qc/0512123)
- [6] Minkevich A V 1980 *Vestsi Akad. Nauk BSSR. Ser. fiz.-mat.*, no. 2 87; 1980 *Phys. Lett. A* **80** 232
- [7] Minkevich A V 1993 *Dokl. Akad. Nauk Belarus.* **37** 33
- [8] Minkevich A V and Garkun A S 2000 *Class. Quantum Grav.* **17** 3045
- [9] Minkevich A V, Garkun A S and Vasilevski Yu G 2004 *Nonlinear Phenomena in Complex Systems* **7** N1 78 (*Preprint* gr-qc/0310060)
- [10] Linde A D 1990 *Particle Physics and Inflationary Cosmology* (Harwood: Chur, Switzerland)
- [11] Bogoliubov N N and Mitropolsky Y A 1961 *Asymptotic Methods in the Theory of Non-Linear Oscillations* (New York: Gordon and Breach Science Publishers)
- [12] Copi C J, Huterer D, Schwarz D J and Starkman G D 2005 On the large-angle anomalies of the microwave sky (*Preprint* astro-ph/0508047)